

SCORE: \_\_\_\_ / 120 POINTS + \_\_\_\_ / 20 POINTS

**NO CALCULATORS ALLOWED****For full credit, you must show the work which leads to all numerical and algebraic answers**Estimate  $\int_{-5}^4 f(x) dx$  using Trapezoidal Rule with  $n = 3$ .

SCORE: \_\_\_\_ / 10 POINTS

|         |     |     |     |      |      |      |      |      |      |     |     |
|---------|-----|-----|-----|------|------|------|------|------|------|-----|-----|
| $x$     | -5  | -4  | -3  | -2   | -1   | 0    | 1    | 2    | 3    | 4   | 5   |
| $f(x)$  | 7   | 9   | 10  | 13   | 12   | 10   | 7    | 3    | 2    | 1   | 5   |
| $f'(x)$ | 1.2 | 0.5 | 1.8 | -0.4 | -1.0 | -2.0 | -3.0 | -1.5 | -0.2 | 2.4 | 1.2 |

$$\Delta x = \frac{4 - (-5)}{3} = 3$$

$$\begin{aligned} T_3 &= \frac{1}{2}(3)(f(-5) + 2f(-2) + 2f(1) + f(4)) \\ &= \frac{3}{2}(7 + 2 \cdot 13 + 2 \cdot 7 + 1) \\ &= \frac{3}{2}(48) = 72 \end{aligned}$$

Give the complete definition of the definite integral.

SCORE: \_\_\_\_ / 10 POINTS

THE DEFINITE INTEGRAL OF  $f$  OVER  $[a, b]$ 

$$\int_a^b f(x) dx = \lim_{n \rightarrow \infty} \sum_{i=1}^n f(x_i) \Delta x \quad \text{WHERE } \Delta x = \frac{b-a}{n}$$

$$\text{AND } a + (i-1)\Delta x \leq x_i \leq a + i\Delta x$$

IF THE LIMIT EXISTS AND IS THE SAME REGARDLESS OF THE CHOICE OF THE  $x_i$ .Compute the exact area under the graph of  $f(x) = 4 - 2x$  on the interval  $[-2, 1]$  using the limit of the right-endpoint Riemann sum.

SCORE: \_\_\_\_ / 20 POINTS

$$\begin{aligned} &\lim_{n \rightarrow \infty} \sum_{i=1}^n f(a + i\Delta x) \Delta x \quad \Delta x = \frac{1 - (-2)}{n} = \frac{3}{n} \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n (4 - 2(-2 + \frac{3i}{n})) \frac{3}{n} \\ &= \lim_{n \rightarrow \infty} \sum_{i=1}^n (8 - \frac{6i}{n}) \frac{3}{n} \\ &= \lim_{n \rightarrow \infty} \frac{3}{n} (8n - \frac{36}{n} \frac{n(n+1)}{2}) \\ &= \lim_{n \rightarrow \infty} 3(8 - \frac{3(n+1)}{n}) \\ &= 3(8 - 3) \\ &= 15 \end{aligned}$$

Find the average value of  $f(x) = \frac{(3x + \sqrt{x})^2}{x^2}$  on  $[1, 4]$ .

SCORE: \_\_\_ / 15 POINTS

$$\begin{aligned} & \frac{1}{4-1} \int_1^4 \frac{(3x + \sqrt{x})^2}{x^2} dx \\ &= \frac{1}{3} \int_1^4 (3 + x^{-\frac{1}{2}})^2 dx \\ &= \frac{1}{3} \int_1^4 (9 + 6x^{-\frac{1}{2}} + x^{-1}) dx \\ &= \frac{1}{3} (9x + 12x^{\frac{1}{2}} + \ln|x|) \Big|_1^4 \\ &= \frac{1}{3} (36 + 24 + \ln 4 - (9 + 12)) \\ &= \frac{1}{3} (39 + \ln 4) \end{aligned}$$

Find the area between  $y = 6x - 3x^2$  and  $y = 0$  on the interval  $[1, 4]$ .

SCORE: \_\_\_ / 15 POINTS

$$\begin{aligned} 6x - 3x^2 &= 0 \\ 3x(2-x) &= 0 \\ x &= 0, 2 \end{aligned}$$

$$\begin{aligned} & \int_1^2 (6x - 3x^2) dx - \int_2^4 (6x - 3x^2) dx \\ &= (3x^2 - x^3) \Big|_1^2 - (3x^2 - x^3) \Big|_2^4 \\ &= (12 - 8) - (3 - 1) - ((48 - 64) - (12 - 8)) \\ &= 4 - 2 - (-16 - 4) \\ &= 22 \end{aligned}$$



Let  $F(x) = \int_3^{3x^3-5x} f(t) dt$ , where  $f(t)$  is the continuous function in the table below. Find  $F'(1)$ .

SCORE: \_\_\_ / 10 POINTS

| $x$     | -5  | -4  | -3  | -2   | -1   | 0    | 1    | 2    | 3    | 4   | 5   |
|---------|-----|-----|-----|------|------|------|------|------|------|-----|-----|
| $f(x)$  | 7   | 9   | 10  | 13   | 12   | 10   | 7    | 3    | 2    | 2   | 5   |
| $f'(x)$ | 1.2 | 0.5 | 1.8 | -0.4 | -1.0 | -2.0 | -3.0 | -1.5 | -0.2 | 2.4 | 1.2 |

$$\begin{aligned} F'(x) &= f(3x^3 - 5x) \cdot \frac{d}{dx}(3x^3 - 5x) \\ &= f(3x^3 - 5x) \cdot (9x^2 - 5) \end{aligned}$$

$$\begin{aligned} F'(1) &= f(-2) \cdot (4) \\ &= 13 \cdot 4 \\ &= 52 \end{aligned}$$

[a]  $\int \frac{4x+1}{\sqrt[3]{3-2x}} dx = -\frac{1}{2} \int (7u^{-\frac{1}{3}} - 2u^{\frac{2}{3}}) du = -\frac{1}{2} \left( \frac{21}{2} u^{\frac{2}{3}} - \frac{6}{5} u^{\frac{5}{3}} \right) + C$   
 $u = 3-2x \rightarrow x = \frac{3-u}{2}$   
 $\frac{du}{dx} = -2$   
 $dx = -\frac{1}{2} du$

$$\frac{4x+1}{\sqrt[3]{3-2x}} dx = \frac{4\left(\frac{3-u}{2}\right)+1}{\sqrt[3]{u}} \cdot -\frac{1}{2} du$$

$$= -\frac{1}{2} \frac{7-2u}{u^{\frac{1}{3}}} du$$

$$= -\frac{1}{2} \int (7u^{-\frac{1}{3}} - 2u^{\frac{2}{3}}) du$$

[b]  $\int_0^1 \frac{6-9x}{3x^2-4x+2} dx = \int_2^1 -\frac{3}{2} \frac{1}{u} du = -\frac{3}{2} \ln|u| = -\left(-\frac{3}{2} \ln 2\right) = \frac{3}{2} \ln 2$   
 $u = 3x^2-4x+2 \rightarrow x=1 \Rightarrow u=1$   
 $x=0 \Rightarrow u=2$   
 $\frac{du}{dx} = 6x-4$   
 $dx = \frac{1}{2(3x-2)} du$

$$\frac{6-9x}{3x^2-4x+2} dx = \frac{-3(3x-2)}{3x^2-4x+2} \cdot \frac{1}{2(3x-2)} du$$

$$= -\frac{3}{2} \cdot \frac{1}{u} du$$

[c]  $\int \frac{6x^3}{1+x^8} dx = \frac{3}{2} \int \frac{1}{1+u^2} du = \frac{3}{2} \tan^{-1} u + C = \frac{3}{2} \tan^{-1} x^4 + C$   
 $u = x^4$   
 $\frac{du}{dx} = 4x^3$   
 $dx = \frac{1}{4x^3} du$

$$\frac{6x^3}{1+x^8} dx = \frac{3 \cancel{6x^3}}{1+x^8} \cdot \frac{1}{\cancel{4x^3} 2} du$$

$$= \frac{3}{2} \frac{1}{1+u^2} du$$

[d]  $\int \left( \sin \frac{x}{3} - \sec 2x \tan 2x + \sec^2 \frac{3x}{2} \right) dx$

$$= -3 \cos \frac{x}{3} - \frac{1}{2} \sec 2x + \frac{2}{3} \tan \frac{3}{2} x + C$$

What month is your birthday? \_\_\_\_\_

What are the first 2 digits of your address? \_\_\_\_\_

What are the last 2 digits of your zip code? \_\_\_\_\_

What are the last 2 digits of your social security number? \_\_\_\_\_

[IF YOU DO NOT HAVE A SOCIAL SECURITY NUMBER,  
USE YOUR STUDENT ID NUMBER]

**You must turn in the no-calculator portion before  
★ you pick up your calculator for this portion ★**

Consider the definite integral  $\int_0^{\frac{\pi}{4}} 3 \cos 2x \, dx$ .

SCORE: \_\_\_\_ / 20 POINTS

- [a] What is the value of  $S_{16}$ ? Round your answer to 4 decimal places.

1.5000

- [b] What is the value of  $M_{20}$ ? Round your answer to 4 decimal places.

1.5003

- [c] Find bounds on the error for the approximation in [b] using the error bounds formula. Show your work.  
You can leave your simplified answer in terms of  $\pi$ .

$$\begin{aligned}
 |EM_{20}| &\leq \frac{K(b-a)^3}{24n^2} \\
 &\leq \frac{12\left(\frac{\pi}{4}-0\right)^3}{12(20)^2} \\
 &= \frac{\frac{\pi^3}{64}}{400} \\
 &= \frac{\pi^3}{25600}
 \end{aligned}$$

$$\begin{aligned}
 f &= 3 \cos 2x \\
 f' &= -6 \sin 2x \\
 f'' &= -12 \cos 2x \\
 |f''| &= 12 \cos 2x \leq 12 \cdot 1 = 12 = K
 \end{aligned}$$