Consider the polar equation $r = 2 - 2\sin 2\theta$.

SCORE: _____ / 16 PTS

POLAR AXIS POLE POLAR AXIS

The following symmetry tests all fail: $(r, -\theta)$, $(-r, \theta)$ and $(-r, \pi - \theta)$

[a] Is the graph symmetric with respect to the polar axis ? **State your conclusion clearly.**

(1) NO CONCLUSION (BOTH TESTS FAIL)

[b] Is the graph symmetric with respect to $\theta = \frac{\pi}{2}$? **State your conclusion clearly.**

$(-r, -\theta)$ $-r = 2 - 2\sin 2(-\theta)$ (½) $(r, \pi - \theta)$

$-r = 2 + 2\sin 2\theta$

$r = -2 - 2\sin 2\theta$ (½)

NO CONCLUSION (1)

$r = 2 - 2\sin 2(\pi - \theta)$ (½)

$r = 2 - 2\sin(2\pi - 2\theta)$

$r = 2 - 2[\sin 2\pi\cos 2\theta - \cos 2\pi \sin 2\theta]$

$r = 2 + 2\sin 2\theta$ (1)

[c] Is the graph symmetric with respect to the pole ? **State your conclusion clearly.**

$(r, \pi + \theta)$ $r = 2 - 2\sin 2(\pi + \theta)$ (½)

$r = 2 - 2\sin(2\pi + 2\theta)$

$r = 2 - 2[\sin 2\pi\cos 2\theta + \cos 2\pi \sin 2\theta]$

(1) $r = 2 - 2\sin 2\theta$ SYMMETRIC (1)

[d] Based on the symmetry tests, what is the minimum interval of the graph you need to plot (before using reflections to draw the rest of the graph) ?

$\theta \in [0, \pi]$ OR $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$ (1) FOR EITHER INTERVAL

[e] Find the zeros of the graph in the minimum interval from [d]
(ie. for what values of θ in the minimum interval does the graph pass through the pole ?).

$0 = 2 - 2\sin 2\theta$

(½) $\sin 2\theta = 1$

$2\theta = \frac{\pi}{2}$ (½) $\theta = \frac{\pi}{4}$ (1)

[f] Find the value of r for all the common values of θ in the minimum interval. Plot those points. Connect the points into a curve. Reflect that part of the curve using the results of the symmetry tests in [a], [b] and [c] to draw the complete graph.



CALCULATE THE r –VALUES ON SCRATCH PAPER ON THE BACK. WRITE THE POLAR COORDINATES HERE.

$(2+\sqrt{3}, -\frac{\pi}{6})$ $(2, 0)$ $(2+\sqrt{3}, \frac{2\pi}{3})$

$(4, -\frac{\pi}{4})$ $(2-\sqrt{3}, \frac{\pi}{6})$ $(4, \frac{3\pi}{4})$

$(2+\sqrt{3}, -\frac{\pi}{3})$ $(0, \frac{\pi}{4})$ $(2+\sqrt{3}, \frac{5\pi}{6})$

$(2, -\frac{\pi}{2})$ $(2-\sqrt{3}, \frac{\pi}{3})$ $(2, \pi)$

$(2, \frac{\pi}{2})$

GRADED BY ME

Consider the conic with polar equation $r = \dfrac{20}{3 - 7\sin\theta}$.

SCORE: ____ / 14 PTS

[a] What is the type of the conic ? **Justify your answer clearly.**

$r = \dfrac{\frac{20}{3}}{1 - \frac{7}{3}\sin\theta}$ $e = \frac{7}{3} > 1$ (1) HYPERBOLA (1)

[b] What is the equation of the directrix ?

$ep = \frac{20}{3}$

$\frac{7}{3}p = \frac{20}{3} \rightarrow p = \frac{20}{7}$ (1) $y = -\frac{20}{7}$ (1) ($\frac{1}{2}$)

[c] Find the **polar AND rectangular** coordinates of the $x-$ and $y-$ intercepts.

θ	r
0	$\frac{20}{3}$
$\frac{\pi}{2}$	-5
π	$\frac{20}{3}$
$\frac{3\pi}{2}$	2

X-INT POLAR $(\frac{20}{3}, 0)$ $(\frac{20}{3}, \pi)$ ($\frac{1}{2}$)

RECT $(\pm\frac{20}{3}, 0)$ ($\frac{1}{2}$)

y-INT POLAR $(-5, \frac{\pi}{2})$ $(2, \frac{3\pi}{2})$ ($\frac{1}{2}$)

RECT $(0, -5)$ $(0, -2)$ ($\frac{1}{2}$)

[d] What are the rectangular coordinates of the vertices, center, foci, and endpoints of the latera recta ?

VERTICES:

$(0, -5)$ $(0, -2)$ (1)

CENTER:

$(0, \frac{-5-2}{2}) = (0, -\frac{7}{2})$ (1)

FOCI:

$(0, 2 \cdot -\frac{7}{2}) = (0, -7)$ (1) AND $(0, 0)$ ($\frac{1}{2}$)

ENDPOINTS OF LATERA RECTA:

$(\pm\frac{20}{3}, 0)$ (1) $(\pm\frac{20}{3}, -7)$ (1)

[e] Graph the conic by connecting the relevant points from [d] appropriately.



GRADED BY ME