Using the distance based definition of a hyperbola, find the equation of the hyperbola with foci  $(0, \pm 12)$ such that the distances from any point on the hyperbola to the foci differ by 4.

SCORE: \_\_\_\_/10 PTS

Do NOT solve this problem using symbols in place of the numbers given.

IF YOU FORGOT ± SEE ALTERNATE SOLUTION IF ORDER

$$\left| \sqrt{x^2 + (y+12)^2} - \sqrt{x^2 + (y-12)^2} \right| = 4$$

OF SUBTRACTION WAS REVERSED

$$\sqrt{x^2 + (y+12)^2} - \sqrt{x^2 + (y-12)^2} = \pm 4$$

 $\sqrt{x^2 + (y+12)^2} = \pm 4 + \sqrt{x^2 + (y-12)^2}$ 

$$48y - 16 = \pm 8\sqrt{x^2 + y^2 - 24y + 144}$$

$$6y - 2 = \pm \sqrt{x^2 + y^2 - 24y + 144}$$

$$36y^2 - 24y + 4 = x^2 + y^2 - 24y + 144$$

$$35y^2 - x^2 = 140$$

$$\frac{y^2}{4} - \frac{x^2}{140} = 1$$

Convert the rectangular equation y = 3x - 5 to polar form. Write r as function of  $\theta$ , and simplify your answer. SCORE: \_\_\_\_\_/4 PTS

 $r\sin\theta = 3r\cos\theta - 5$ 

$$r\sin\theta - 3r\cos\theta = -5$$

$$r(\sin\theta - 3\cos\theta) = -5$$

$$r = \frac{-5}{\sin \theta - 3\cos \theta} = \frac{5}{3\cos \theta - \sin \theta}$$

Fill	in	the	blanks
1 111	ш	uic	Ulains.

SCORE: /7 PTS

A house has an exposed (straight) beam 25 feet above and parallel to the floor. A small lamp hangs from the ceiling. [a]

There is an arch such that the distance from any point on the arch to the lamp is the same as the distance from that point to the beam.

- The shape of the arch is a/an PARABOLA (or part of it).

  The shape of the graph of the equation  $3x^2 3x + 2y^2 2y 1 = 0$  is a/an PARABOLA. [6]
- The shape of the graph of the equation  $3x^2 + 2x + 3y^2 3y 1 = 0$  is a/an  $\frac{1}{1}$ [c]
- The polar co-ordinates  $(-5, \frac{4\pi}{7})$  refer to the same point as the polar co-ordinates  $(5, \frac{1177}{7})$ . (Your answer must be **positive**.) [d]
- The polar co-ordinates  $(5, \frac{4\pi}{7})$  refer to the same point as the polar co-ordinates  $(5, \frac{10\pi}{7})$ . (Your answer must be <u>negative</u>.) [e]
- The point with polar co-ordinates  $(-5, -\frac{2\pi}{3})$  lies in quadrant [f]

Convert the polar equation  $r^2 = \cos 2\theta$  to rectangular form.

SCORE: /4 PTS

GRADE AGAINST

VERSION OF

SOLUTION

Simplify your answer so that there are no radicals, complex fractions, fractional exponents nor negative exponents.

$$r^{2} = \cos^{2}\theta - \sin^{2}\theta$$

$$r^{2} = \left(\frac{x}{r}\right)^{2} - \left(\frac{y}{r}\right)^{2}$$

$$r^2 = \cos^2 \theta - \sin^2 \theta$$

$$r^2r^2 = r^2(\cos^2\theta - \sin^2\theta)$$

$$r^{2} = \frac{x^{2}}{r^{2}} - \frac{y^{2}}{r^{2}}$$

$$r^{4} = x^{2} - y^{2}$$

$$(x^{2} + y^{2})^{2} = x^{2} - y^{2}$$

$$r^4 = r^2 \cos^2 \theta - r^2 \sin^2 \theta$$

$$r^4 = r^2 - v^2$$

Find the foci, and equations of the asymptotes of the hyperbola  $x^2 - 2y^2 - 4x - 16y - 22 = 0$ .

SCORE: \_\_\_\_\_/ 5 PTS

$$(x^2 - 4x) - 2(y^2 + 8y) = 22$$

$$(x^2 - 4x + 4) - 2(y^2 + 8y + 16) = 22 + 4 - 32$$

$$(x^{2}-4x+4)-2(y^{2}+8y+16) = 22+4-32$$


$$(x-2)^{2}-2(y+4)^{2}=-6$$

$$\frac{(y+4)^{2}}{3}-\frac{(x-2)^{2}}{6}=1$$

$$c^2 = 3 + 6 = 9 \implies c = 3$$

$$c^2 = 3 + 6 = 9 \implies c = 3$$
  
slopes of asymptotes  $= \pm \sqrt{\frac{3}{6}} = \pm \sqrt{\frac{1}{2}} = \pm \frac{\sqrt{2}}{2}$ 

$$(2, -4 \pm 3) = (2, -7), (2, -1)$$



## TERNATE SOLUTION ONLY (1) IF YOU FORGOT + $\left| \sqrt{x^2 + (y - 12)^2} - \sqrt{x^2 + (y + 12)^2} \right| = 4$ $\sqrt{x^2 + (y-12)^2} - \sqrt{x^2 + (y+12)^2} = \pm 4$ $\sqrt{x^2 + (y-12)^2} = \pm 4 + \sqrt{x^2 + (y+12)^2}$ $(x^2 + y^2 - 24y + 144) = (16 \pm 8\sqrt{x^2 + y^2 + 24y + 144} + x^2 + y^2 + 24y + 144)$ $-48y - 16 = \pm 8\sqrt{x^2 + y^2 + 24y + 144}$

$$-48y - 16 = \pm 8\sqrt{x^2 + y^2 + 24y + 144}$$
$$-6y - 2 = \pm \sqrt{x^2 + y^2 + 24y + 144}$$

 $35y^2 - x^2 = 140$ 

 $36y^2 + 24y + 4 = x^2 + y^2 + 24y + 144$