

$$\text{Find } \frac{d}{dx}(\tanh^{-1} x^2 + \operatorname{sech} 3x).$$

SCORE: _____ / 15 PTS

You may use any hyperbolic identities or the derivatives of any hyperbolic functions without proving them.

$$\begin{aligned} & \frac{1}{1-(x^2)^2} \cdot 2x + (-\operatorname{sech} 3x \tanh 3x)(3) \\ \textcircled{5} \quad &= \frac{2x}{1-x^4} - 3\operatorname{sech} 3x \tanh 3x \end{aligned}$$

③ ④ ③

The graph of $f(t)$ is shown on the right. **NOTE: The graph consists of a quarter circle, a semi-circle, a line segment, another quarter circle, and another line segment.**

SCORE: ____ / 20 PTS

Let $g(x) = \int_5^x f(t) dt$.

- [a] Find $g(-2)$.

$$\int_5^2 f(t) dt = - \int_{-2}^5 f(t) dt = - \left[\frac{1}{4}\pi(2)^2 - \frac{1}{2}(3)(3) - \frac{1}{4}\pi(2)^2 - 2 \right] = \frac{13}{2}$$

- [b] Find $g'(-4)$.

$$f(-4) = \boxed{\textcircled{1}} \quad \boxed{\textcircled{2}}$$


- [c] Find $g''(2)$.

$$f'(2) = -1 \quad \boxed{\textcircled{2}}$$

- [d] Find the x -coordinates of all local maxima of g .

$g'(x) = f(x)$ CHANGES FROM POSITIVE TO NEGATIVE $\boxed{\textcircled{3}}$

$\boxed{\textcircled{2}} \quad x = 0 \quad \boxed{\textcircled{3}}$



Prove that $\frac{\pi}{2} \leq \int_1^{\sqrt{3}} x^3 \tan^{-1} x \, dx \leq \frac{2\pi}{3}$.

SCORE: _____ / 15 PTS

ON $[1, \sqrt{3}]$, (5) $\frac{\pi}{4} \leq \tan^{-1} x \leq \frac{\pi}{3}$

$$\frac{\pi}{4}x^3 \leq x^3 \tan^{-1} x \leq \frac{\pi}{3}x^3$$

$$\begin{aligned} & \int_1^{\sqrt{3}} x^3 \, dx \\ (2) &= \left[\frac{1}{4}x^4 \right]_1^{\sqrt{3}} \\ &= \frac{1}{4}(9-1) = 2 \end{aligned}$$

(4) $\int_1^{\sqrt{3}} \frac{\pi}{4}x^3 \, dx \leq \int_1^{\sqrt{3}} x^3 \tan^{-1} x \, dx \leq \int_1^{\sqrt{3}} \frac{\pi}{3}x^3 \, dx$

$$\frac{\pi}{2} = 2\left(\frac{\pi}{4}\right) \leq \int_1^{\sqrt{3}} x^3 \tan^{-1} x \, dx \leq 2\left(\frac{\pi}{3}\right) = \frac{2\pi}{3}$$

(4)

Find $\lim_{n \rightarrow \infty} \frac{4}{n} \sum_{i=1}^n \frac{1}{3 + \frac{4i}{n}}$ by finding the corresponding definite integral, and evaluating that integral.

SCORE: _____ / 15 PTS

$$\begin{aligned} & \text{③ } \int_3^7 \frac{1}{x} dx = \left[\ln|x| \right]_3^7 \\ & \text{③ } \int_3^7 \frac{1}{x} dx = \left[\ln|x| \right]_3^7 \\ & = \ln 7 - \ln 3 \\ & = \ln \frac{7}{3} \quad \text{③} \end{aligned}$$

$$a + i\Delta x = 3 + \frac{4i}{n}$$

$$a = 3, \Delta x = \frac{4}{n} = \frac{b-a}{n}$$

$$\frac{4}{n} = \frac{b-3}{n}$$

$$b = 7$$

$$f(a + i\Delta x) = f\left(3 + \frac{4i}{n}\right) = \frac{1}{3 + \frac{4i}{n}}$$

$$f(x) = \frac{1}{x}$$

The velocity of a car at time t hours after 1pm is given by $v(t) = t^2 - 3t - 4$ miles per hour.

SCORE: _____ / 15 PTS

Find the total distance travelled by, NOT THE DISPLACEMENT OF, the car from 2 pm to 5 pm.

$$\int_1^4 |v(t)| dt = \int_1^4 |t^2 - 3t - 4| dt$$


$$t=1 \quad t=4$$

$$\begin{aligned} &= \int_1^4 (-t^2 + 3t + 4) dt \\ &\quad \text{①} \quad \text{⑤} \\ &= \left[-\frac{1}{3}t^3 + \frac{3}{2}t^2 + 4t \right]_1^4 \\ &\quad \text{④} \end{aligned}$$

$$= -\frac{1}{3}(64-1) + \frac{3}{2}(16-1) + 4(4-1)$$

$$= -21 + \frac{45}{2} + 12 = \frac{27}{2} \text{ miles} \quad \text{①}$$

$$\begin{aligned} &t^2 - 3t - 4 \\ &= (t-4)(t+1) \end{aligned}$$



Evaluate the following integrals. If a definite integral cannot be evaluated using the Fundamental Theorem of Calculus, write "FTC DOES NOT APPLY", and explain why not with proper justification.

SCORE: _____ / 55 PTS

[a] $\int_{-1}^2 \frac{1-x}{\sqrt{5+6x}} dx$

(3) FTC DOES NOT APPLY -
 (3) DISCONTINUOUS (DOES NOT EXIST) @ $x = -\frac{5}{6}$

[b] $\int_{-3}^3 \frac{7t^2}{(4-t^2)^5} dt$

(3) FTC DOES NOT APPLY -
 (3) DISCONTINUOUS @ $t = \pm 2$

[c] $\int \frac{5e^{2y}}{\sqrt{1-e^{4y}}} dy$

(4) $u = e^{2y}$
 $du = 2e^{2y} dy$
 $\frac{5}{2} du = 5e^{2y} dy$

(5) $\int \frac{\frac{5}{2} du}{\sqrt{1-u^2}}$

(5) $= \frac{5}{2} \sin^{-1} u + C$

(4) $= \frac{5}{2} \sin^{-1} e^{2y} + C$ (2)

[d] $\int_{-\pi}^{\pi} x^5 \cos x dx$

(4) $(-x)^5 \cos(-x) = -x^5 \cos x$
 (4) ODD, CONTINUOUS (3)
INTEGRAL = 0

(4)