Math 43 Midterm 1 Review

HYPERBOLICS

REFER TO THE HYPERBOLIC FUNCTIONS SUPPLEMENT

POLAR

[1] Remember that a single point in the plane has infinitely many polar co-ordinates. Consider the point with polar co-ordinates $(7, \frac{2\pi}{3})$.

- [a] Find another pair of polar co-ordinates for this point, using a positive r value, and a positive θ value.
- [b] Find another pair of polar co-ordinates for this point, using a positive r value, and a negative θ value.
- [c] Find another pair of polar co-ordinates for this point, using a negative r value, and a positive θ value.
- [d] Find another pair of polar co-ordinates for this point, using a negative r value, and a negative θ value.
- [2] Convert the following points or equations.
 - [a] the point with polar co-ordinates $(8, \frac{5\pi}{6})$ to rectangular co-ordinates
 - [b] the point with rectangular co-ordinates (-6, -2) to polar co-ordinates
 - [c] the rectangular equation $x^2 y^2 2x = 0$ to polar
 - [d] the polar equation $r = \frac{7}{4 2\cos\theta}$ to rectangular
 - [e] the rectangular equation 3x 2y + 6 = 0 to polar
 - [f] the polar equation $r = \cos 2\theta$ to rectangular
 - [g] the rectangular equation $x^2 + 6y 9 = 0$ to polar
 - [h] the polar equation $\theta = \frac{5\pi}{6}$ to rectangular
- [3] Run the standard tests for symmetry for the polar equation $r^3 = 1 \sin 2\theta$, and state the conclusions. What is the minimum interval of θ -values that must be plotted before using symmetry to complete the graph?
- [4] Find the values of $\theta \in [0, 2\pi)$ at which the graph of the polar equation $r = 2\cos 2\theta + 1$ passes through the pole.
- [5] Name the shape of the graphs of the following polar equations. If the graph is a rose curve, state the number of petals.
 - r = 5 $r = 6 4\cos\theta$ $r = 5 - 5\sin\theta$ $r = 7 \sin 6\theta$ $\theta = 7$ [c] [d] [a] [b] $r = 2 + 3\cos\theta$ $r = 3\sin\theta$ $r = 4 \sin 9\theta$ [f] [g] [h] [e] $r = 6 + 2\sin\theta$ [i]

[6] Determine if each polar equation corresponds to a circle, a parabola, an ellipse or a hyperbola. If the equation corresponds to a circle, find its center & radius.

If the equation corresponds to a parabola, find its eccentricity, focus, directrix & vertex.

If the equation corresponds to an ellipse, find its eccentricity, foci, directrix, center & the endpoints of the major axes and latera recta. If the equation corresponds to a hyperbola, find its eccentricity, foci, directrix, center, vertices & the endpoints of the latera recta. **Do not convert the equations to rectangular co-ordinates.**

Final answers must be in rectangular co-ordinates.

[a]
$$r = \frac{10}{3 - 3\sin\theta}$$
 [b] $r = \frac{10}{3 - 2\cos\theta}$ [c] $r = \frac{10}{2 + 3\sin\theta}$ [d] $r = 10$

- [7] Find the polar equations of the following conics with their focus at the pole.
 - [a] Parabola: directrix x = 7
 - [b] Parabola: vertex $(7, \frac{3\pi}{2})$
 - [c] Ellipse: eccentricity $\frac{3}{4}$, directrix y = 5
 - [d] Ellipse: vertices (4, 0) and $(2, \pi)$
 - [e] Hyperbola: eccentricity $\frac{5}{2}$, directrix x = -3
 - [f] Hyperbola: vertices $(3, \frac{3\pi}{2})$ and $(15, \frac{3\pi}{2})$
- [8] Draw diagrams and write algebraic equations involving distances to answer the following questions.

A drinking fountain is 15 feet from the wall of a school building.

- [a] A cat is running on the school grounds, so that it is always three times as far from the wall as it is from the fountain. What is the shape of the cat's path ?
- [b] A dog is running on the school grounds, so that it is always three times as far from the fountain as it is from the wall. What is the shape of the dog's path ?
- [c] A chicken is running on the school grounds, so that it is always as far from the wall as it is from the fountain. What is the shape of the chicken's path ?
- [9] Sketch the graphs of the polar equations in [5][a], [f], [g] and [i] using the shortcut process shown in lecture. Find all x - and y - intercepts.

ANSWERS

POLAR

- [1] [a] $(7, \frac{8\pi}{3})$ [b] $(7, -\frac{4\pi}{3})$ [c] $(-7, \frac{5\pi}{3})$ [d] $(-7, -\frac{\pi}{3})$
- [2] [a] $(-4\sqrt{3}, 4)$ [b] $(2\sqrt{10}, 3.46)$ $2\cos\theta$

[c]
$$r = \frac{2\cos\theta}{\cos 2\theta} = 2\cos\theta \sec 2\theta$$
 [d] $12x^2 + 16y^2 - 28x - 49 = 0$
[e] $r = \frac{6}{10x^2 + 10x^2}$ [f] $(x^2 + y^2)^3 = (x^2 - y^2)^2$

[e]
$$r = \frac{1}{2\sin\theta - 3\cos\theta}$$

[g] $r = \frac{3}{1 + \sin\theta} or \frac{-3}{1 - \sin\theta}$
[f] $(x^2 + y^2)^3 = (x^2 - y^2)^3$
[h] $y = -\frac{\sqrt{3}}{3}x$

[3]	Symmetry over polar axis	substituting $(r, -\theta)$	gives $r^3 = 1 + \sin 2\theta$	no conclusion
		substituting $(-r, \pi - \theta)$	gives $r^3 = -1 - \sin 2\theta$	no conclusion
	Symmetry over pole:	substituting $(-r, \theta)$	gives $r^3 = -1 + \sin 2\theta$	no conclusion
		substituting $(r, \pi + \theta)$	gives $r^3 = 1 - \sin 2\theta$	symmetric over pole
	Symmetry over $\theta = \frac{\pi}{2}$:	substituting $(-r, -\theta)$	gives $r^3 = -1 - \sin 2\theta$	no conclusion
		substituting $(r, \pi - \theta)$	gives $r^3 = 1 + \sin 2\theta$	no conclusion

Minimum interval $\theta \in [0, \pi]$ or $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$

rose curve with 9 petals

convex limacon

[f]

[e]

[i]

[4]
$$0 = 2\cos 2\theta + 1 \text{ for } 0 \le \theta < 2\pi$$
$$\cos 2\theta = -\frac{1}{2}$$
$$2\theta = \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{8\pi}{3}, \frac{10\pi}{3} \text{ since } 0 \le 2\theta < 4\pi$$
$$\theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$$
[5] [a] cardioid [b] rose curve with 12 petals [c] circle [d] line

limacon with inner loop

[g]

limacon with dimple

[h]

circle

[6]	[a]	PARABOLA Eccentricity: Focus:	1 (0, 0)		
		Directrix:	$y = -\frac{10}{3}$		
		Vertex:	$(0, -\frac{5}{3})$		
	[b]	ELLIPSE			
		Eccentricity:	$\frac{2}{3}$		
		Foci:	(0,0) and $(8,0)$		
		Directrix:	x = -5		
		Center:	(4, 0)		
		Endpoints of major axis:	(-2, 0) and $(10, 0)$		
		Endpoints of latera recta:	$(0,\pm\frac{10}{3})$ and $(8,\pm\frac{10}{3})$		
	[_]				
	[c]	HYPERBOLA Eccentricity:	$\frac{3}{2}$		
		Foci:	(0, 0) and $(0, 12)$		
		Directrix:	$y = \frac{10}{3}$		
		Center:	$y = \frac{1}{3}$ (0, 6)		
		Vertices:	(0, 0) (0, 2) and $(0, 10)$		
		Endpoints of latera recta:			
		Enupoints of fatera feeta.	$(\pm 5, 0)$ and $(\pm 5, 12)$		
	[d]	Center:	(0,0)		
		Radius:	10		
		7	14	15	
[7]	[a]	$r = \frac{7}{1 + \cos\theta}$	[b] $r = \frac{14}{1 - \sin \theta}$	[c] $r = \frac{15}{4+3\sin\theta}$	
		$1 + \cos\theta$	$1 - \sin \theta$	$4 + 5 \sin \theta$	
	[d]	$r = \frac{6}{3 - \cos\theta}$	$[e] \qquad r = \frac{15}{2 - 5\cos\theta}$	[f] $r = \frac{15}{2 - 3\sin\theta}$	
[8]	[a]	ellipse	[b] (part of) hyperbola	[c] parabola	
[9]	[a]	$r = 5 - 5\sin\theta$	[f] $r = 2 + 3\cos\theta$	[g] $r = 6 - 4\cos\theta$	[i] $r = 6 + 2\sin\theta$
		8	8- 8-	8	
		6+ 4-	6+ 4		
		2-	·····		
	-* -	4 -2 -2 4 6 8 -2 -		-8 -6 -4 -2 4 6 8 -2-	
			-6		-6
		-8	-8	-8 -	
	(±5, 0),	(5,0), (1,0), (0,0),	(2,0), (-10,0),	$(\pm 6, 0),$
		, (0, -10)	$(0, \pm 2)$	$(0, \pm 6)$	(0, 8), (0, -4)
					· ·