

If $p(x) = \int_{4x}^{x^3} \tan^{-1} e^{2t} dt$, find $p'(x)$.

SCORE: ____ / 4 PTS

$$p(x) = \int_{4x}^0 \tan^{-1} e^{2t} dt + \int_0^{x^3} \tan^{-1} e^{2t} dt$$
$$= -\int_0^{4x} \tan^{-1} e^{2t} dt + \int_0^{x^3} \tan^{-1} e^{2t} dt$$

$$p'(x) = \frac{d}{dx} \int_0^{4x} \tan^{-1} e^{2t} dt + \frac{d}{dx} \int_0^{x^3} \tan^{-1} e^{2t} dt$$

$$= -\frac{d}{d(4x)} \int_0^{4x} \tan^{-1} e^{2t} dt \cdot \frac{d(4x)}{dx} + \frac{d}{d(x^3)} \int_0^{x^3} \tan^{-1} e^{2t} dt \cdot \frac{d(x^3)}{dx}$$


$$= -\tan^{-1} e^{8x} \cdot 4 + \tan^{-1} e^{2x^3} \cdot 3x^2$$

$$= \boxed{3x^2 \tan^{-1} e^{2x^3}} - \boxed{4 \tan^{-1} e^{8x}} \quad \text{ALL ITEMS WORTH 1}$$

Let $g(x) = \int_{-3}^x f(t) dt$, where f is the function whose graph is shown on the right.

SCORE: ____ / 7 PTS

- [a] Write "I UNDERSTAND THAT THE GRAPH SHOWS f , BUT THE QUESTIONS ASK ABOUT g ".



- [b] Find $g'(1)$. Explain your answer very briefly.

$$g'(1) = \underbrace{f(1)} = -1$$

- [c] Find all critical numbers of g . Explain your answer very briefly.

$$g'(x) = \underbrace{f(x)} = 0 \text{ @ } \underbrace{x = -5, 2}$$

(2)

ALL ITEMS WORTH (1)
EXCEPT AS INDICATED

- [d] Find all intervals over which g is both decreasing and concave up at the same time. Explain your answer very briefly.

$$g'(x) = \underbrace{f(x)} < 0 \text{ AND } \underbrace{\text{INCREASING}} \text{ ON } \underbrace{(-1, 2)}$$

Evaluate the following integrals, or explain why they can't be evaluated.

SCORE: ____ / 11 PTS

[a] $\int_0^{\frac{\pi}{2}} \frac{\sin r \cos r}{1 + \cos^4 r} dr$

[b] $\int_{-3}^3 \frac{t^3}{1-t^6} dt$

$u = \cos^2 r$ ①
 $du = -2 \cos r \sin r dr$
 $-\frac{1}{2} du = \cos r \sin r dr$

DISCONTINUOUS @ $t = \pm 1$ ①
 FTC DOESN'T APPLY

$r = 0 \rightarrow u = 1$
 $r = \frac{\pi}{2} \rightarrow u = 0$

① $\int_1^0 -\frac{1}{2} \frac{1}{1+u^2} du$ ①
 $= -\frac{1}{2} \tan^{-1} u \Big|_1^0$
 $= -\frac{1}{2} (0 - \frac{\pi}{4})$
 $= \frac{\pi}{8}$

ALL ITEMS WORTH $\left(\frac{1}{2}\right)$
 EXCEPT AS INDICATED

[c] $\int \frac{(2\sqrt{y} - 3y^2)^2}{6y^5} dy$

[d] $\int_{-\pi}^{\pi} \frac{\sin^3 \theta}{\sqrt{4 + \cos \theta}} d\theta = 0$

$\int \frac{4y - 12y^{\frac{5}{2}} + 9y^4}{6y^5} dy$

$\frac{\sin^3(-\theta)}{\sqrt{4 + \cos(-\theta)}} = -\frac{\sin^3 \theta}{\sqrt{4 + \cos \theta}}$ ①

$= \int \left(\frac{2}{3}y^{-4} - 2y^{-\frac{5}{2}} + \frac{3}{2}y^{-1}\right) dy$ ①

ODD, CONTINUOUS

$= \frac{2}{3} \left(-\frac{1}{3}\right) y^{-3} - 2 \left(-\frac{2}{3}\right) y^{-\frac{3}{2}} + \frac{3}{2} \ln|y| + C$

$= -\frac{2}{9} y^{-3} + \frac{4}{3} y^{-\frac{3}{2}} + \frac{3}{2} \ln|y| + C$

In complete sentences, using proper English and mathematical notation,

state the Fundamental Theorem of Calculus (both parts) and the Net Change Theorem.

SCORE: ____ / 5 PTS

IF f IS CONTINUOUS ON $[a, b]$

① IF $g(x) = \int_a^x f(t) dt$, THEN $g'(x) = f(x)$

② IF $F'(x) = f(x)$, THEN $\int_a^b f(t) dt = F(b) - F(a)$

IF F' IS CONTINUOUS ON $[a, b]$

THEN $\int_a^b F'(x) dx = F(b) - F(a)$

GRADED BY ME

If $f(a)$ is the annual amount of weight Morgan gained (in kilograms per year) when she was a years old,

SCORE: ____ / 3 PTS

what is the meaning of the equation $\int_{14}^{16} f(x) dx = 12$?

NOTES: Your answer must use all three numbers from the equation, along with correct units.
Your answer should NOT use “a”, “x”, “f(x)”, “integral”, “antiderivative”, “rate of change” or “derivative”.
Your answer should sound like normal spoken English.

FROM AGE 14 YEARS OLD TO 16 YEARS OLD,
MORGAN'S WEIGHT INCREASED 12 kg ALTOGETHER

GRADED BY ME