## Math 1A Midterm 1 Review

- IF YOU HAVE TAKEN DIFFERENTIAL CALCULUS BEFORE, DO NOT USE DIFFERENTIATION SHORTCUTS.
- YOU SHOULD ONLY REQUIRE A CALCULATOR FOR QUESTIONS MARKED [C].
- UNLESS A GRAPH IS GIVEN, YOU MUST BE ABLE TO SOLVE EACH PROBLEM WITHOUT A GRAPH.
- Estimate the slope of the tangent line to the curve  $y = \sqrt{x + \sqrt{\cos x}}$  at the point (0, 1) using the slopes of several secant lines.
- The position of an object (in meters) at time t seconds, is given by the function  $f(t) = t^2 \cos \pi t$ . Find the average velocity of the [2] object over the interval [1, 5]. Specify the units.
- [3] Sketch the graph of a function f(x) which satisfies the following conditions:

$$\lim_{x \to -2^+} g(x) = -3, \qquad \lim_{x \to -2^-} g(x) = \infty, \qquad \lim_{x \to 1} g(x) = -\infty, \qquad \lim_{x \to -\infty} g(x) = 2, \text{ and} \qquad \lim_{x \to \infty} g(x) = -2$$
Your graph should be continuous at all points unless otherwise required by the conditions above.

Prove that  $\lim_{x\to 0} x^4 \cos \frac{1}{x^2} = 0$ . [4]

[5] Let 
$$f(x) = \begin{cases} 2x-3 & \text{if } x < -1 \\ x^2 - 6 & \text{if } -1 < x < 2 \\ 4x - 6 & \text{if } x \ge 2 \end{cases}$$

[a] Find 
$$\lim_{x \to -2} f(x)$$
.

[b] Find 
$$\lim_{x \to -1} f(x)$$
.  
[c] Find  $\lim_{x \to 2} f(x)$ .

[c] Find 
$$\lim_{x\to 2} f(x)$$

- Find the value of a if  $\lim_{x\to 2} \frac{\sqrt{x^2 + a} 1}{x 2} = 2$ . [6]
- If  $\lim_{x\to 2} f(x) = -3$  and  $\lim_{x\to 2} g(x) = 4$ , find  $\lim_{x\to 2} \frac{x^2 g(x)}{1+f(x)}$ . Show clearly how the limit laws are used in your solution. [7]
- Find the discontinuities of  $f(x) = \frac{x+2}{x^2-9}$ , and find the one-sided limits at each discontinuity. [8]

[9] Let 
$$f(x) = \begin{cases} 2x + a & \text{if } x < -1 \\ 3 - x & \text{if } -1 < x < 2 \\ bx - 1 & \text{if } x \ge 2 \end{cases}$$

- [a] Find the value of a so that f(x) is continuous at x = -1.
- Find the value of b so that f(x) is continuous at x = 2. [b]
- If a = 6 and b = 3, find all discontinuities of f(x) and find the type of each discontinuity (removable, jump or infinite). [c]
- Use the Intermediate Value Theorem to prove that the equation  $\cos 2x = x^2$  has a solution in the interval  $[0, \pi]$ . [10]

- Find all horizontal and vertical asymptotes of  $f(x) = \frac{\sqrt{4+9x^2}}{2x-1}$ . [11]
- If  $f(x) = x^3 3x + 2$ , find f'(-2) using both definitions of f'(a). [12]
- Find a function f and a number a such that the derivative of f at a is given by [13]
  - $\lim_{h\to 0} \frac{\cos(\pi(h-1))+1}{h}$ [a]


- $\lim_{x \to -2} \frac{x^2 x 6}{x + 2}$ [b]
- The position of an object (in feet) at time t minutes, is given by the function  $f(t) = \sqrt{t^2 5}$ . Find the instantaneous velocity of [14] the object at time t = 3. Specify the units.
- Find the equation of the tangent line to the curve of  $f(x) = \frac{2x}{1-x}$  at x=2. [15]
- [16] The graph of f is shown to the right. Arrange the following from least (most negative) to greatest (most positive).



$$f'(-4)$$

$$f'(-2)$$





The time required to defrost a package of frozen food in the refrigerator depends on the temperature inside the refrigerator. [17]


Let t = f(T), where t is the defrost time (in hours), and T is the refrigerator temperature (in  ${}^{\circ}C$ )

- Give the practical meaning (including units) of f(4) = 6. [a]
- Give the practical meaning (including units) of f'(4) = -1. [b]
- Is there a value of  $T_0$  for which you would expect  $f'(T_0) > 0$  ? Why or why not ? [c]
- Using the definition of the derivative, find the derivatives of the following functions. [18]

[a] 
$$f(t) = \frac{1}{\sqrt{1-t}}$$

$$[b] g(x) = \frac{4x}{2-x}$$

- The graph of f(x) is shown on the right. [19]
  - Find all x-coordinates where f'(x) is undefined, [a] and explain briefly why.
  - Sketch a graph of f'(x). [b]



If the tangent line to the graph of y = f(x) at x = 4 is x - 2y = 6, prove that  $\lim_{x \to 4} f(x) = -1$ .

## YOU MUST ALSO KNOW THE FOLLOWING DEFINITIONS AND THEOREMS:

**Definitions** 

[20]

vertical/horizontal asymptote (from textbook) continuity at a point (from lecture) removable discontinuity (from lecture) jump discontinuity (from lecture) derivative at a point derivative function

## Theorems

Squeeze Theorem Intermediate Value Theorem Differentiability implies continuity