


Consider the graph of the polar equation $r = 8 - 5 \sin \theta$.

SCORE: ____ / 6 PTS



[a] Fill in the blanks.

[i] The shape of the graph is a/an LIMACON WITH DIMPLE. ①

[ii] The graph DOES NOT pass through the pole.
does / does not

[iii] Find the rectangular coordinates of the

x - intercept(s)

① $(\pm 8, 0)$

y - intercept(s)

$(0, 3)$ $(0, 13)$

② $\frac{1}{2}$

③ $\frac{1}{2}$

[b] Sketch the graph on the grid provided above. You must provide a scale for the axes & plot all points from part [a][iii] above.

Fill in the blanks.

SCORE: ____ / 4 PTS

- [a] The polar co-ordinates $(-11, -\frac{8\pi}{5})$ correspond to the same point as the polar co-ordinates $(-11, \frac{2\pi}{5})$ and $(11, \frac{7\pi}{5})$.

NOTE: Both your answers for this question must be positive.

- [b] The point with rectangular co-ordinates $(-3\sqrt{2}, -\sqrt{6})$ has polar co-ordinates $(2\sqrt{6}, \frac{7\pi}{6})$.

NOTE: Both parts of your answer for this question must be positive.

- [c] The point with polar co-ordinates $(8, -\frac{2\pi}{3})$ has rectangular co-ordinates $(-4, -4\sqrt{3})$.

$$x = 8 \cos(-\frac{2\pi}{3}) = 8 \cdot -\frac{1}{2}$$

$$y = 8 \sin(-\frac{2\pi}{3}) = 8 \cdot -\frac{\sqrt{3}}{2}$$

$$r^2 = (-3\sqrt{2})^2 + (-\sqrt{6})^2 = 18 + 6 = 24$$
$$\theta = \pi + \tan^{-1} \frac{-\sqrt{6}}{-3\sqrt{2}} = \pi + \tan^{-1} \frac{\sqrt{3}}{3} = \pi + \frac{\pi}{6}$$

Convert the polar equation $r = 2 - 7 \sin 2\theta$ to rectangular, and simplify as shown in the website handout. SCORE: ____ / 5 PTS

NOTE: Your final answer should **NOT** have fractions, fractional exponents nor radicals.

$$r = 2 - 7(2 \sin \theta \cos \theta)$$

$$r = 2 - 14 \frac{y}{r} \frac{x}{r}$$

$$r^3 = 2r^2 - 14xy$$

$$(\sqrt{x^2 + y^2})^3 = 2(x^2 + y^2) - 14xy$$
$$= 2x^2 - 14xy + 2y^2$$

$$(x^2 + y^2)^3 = (2x^2 - 14xy + 2y^2)^2$$

① POINT EACH

Consider the graph of the polar equation $r = 4 - 8\cos 4\theta$.

SCORE: ____ / 15 PTS

NOTE: $(-r, \theta)$ and $(-r, \pi - \theta)$ tests do NOT show that the graph is symmetric

POLE POLAR AXIS

- [a] Using the information above, and the tests and shortcuts shown in lecture, test if the graph is symmetric over the pole, the polar axis, and/or $\theta = \frac{\pi}{2}$. State your conclusions in the table. **NOTE: Run as FEW tests as needed to prove your answers are correct.**

POLE: $r = 4 - 8\cos 4(\pi + \theta)$ ①
 $r = 4 - 8\cos (4\pi + 4\theta)$
 $r = 4 - 8[\cos 4\pi \cos 4\theta - \sin 4\pi \sin 4\theta]$

① $r = 4 - 8\cos 4\theta$, SYM

POLAR AXIS: $r = 4 - 8\cos 4(-\theta)$ ①
 $r = 4 - 8\cos 4\theta$, SYM

① $r = 4 - 8\cos 4\theta$, SYM

AUTOMATICALLY SYM OVER $\theta = \frac{\pi}{2}$

Type of symmetry	Conclusion
Over the polar axis	SYMMETRIC
Over $\theta = \frac{\pi}{2}$	SYMMETRIC
Over the pole	SYMMETRIC

① POINT IF EXACTLY 2 CORRECT
 ② POINTS IF ALL CORRECT

- [b] Based on the results of part [a], what is the minimum interval of the graph you need to plot first (before using reflections to draw the rest of the graph)?

$[0, \frac{\pi}{2}]$ ①

- [c] Find the angles algebraically in the minimum interval in part [b] at which the graph goes through the pole.


$0 = 4 - 8\cos 4\theta$
 $8\cos 4\theta = 4$
 $\cos 4\theta = \frac{1}{2}$ ①
 $0 \leq \theta \leq \frac{\pi}{2} \rightarrow 0 \leq 4\theta \leq 2\pi$
 $4\theta = \frac{\pi}{3}, \frac{5\pi}{3}$ ①
 $\theta = \frac{\pi}{12}, \frac{5\pi}{12}$ ①

- [d] Find the value of r (rounded to 1 decimal place) for all common angles in the minimum interval in part [b].

NOTE: You do NOT need to show work, only answers.

θ	r
0	-4
$\frac{\pi}{6}$	8
$\frac{\pi}{4}$	12
$\frac{\pi}{3}$	8
$\frac{\pi}{2}$	-4

① POINT EACH



GRADED BY ME

- [d] Sketch the graph on the grid provided below. **You must provide a scale for the polar axis & plot all points from part [c] above.**