

Math 43 Midterm 1 Review

HYPERBOLICS

REFER TO THE HYPERBOLIC FUNCTIONS SUPPLEMENT

POLAR


- [1] Remember that a single point in the plane has infinitely many polar co-ordinates.
Consider the point with polar co-ordinates $(7, \frac{2\pi}{3})$.
- [a] Find another pair of polar co-ordinates for this point, using a positive r – value, and a positive θ – value.
[b] Find another pair of polar co-ordinates for this point, using a positive r – value, and a negative θ – value.
[c] Find another pair of polar co-ordinates for this point, using a negative r – value, and a positive θ – value.
[d] Find another pair of polar co-ordinates for this point, using a negative r – value, and a negative θ – value.
- [2] Convert the following points or equations.
- [a] the point with polar co-ordinates $(8, \frac{5\pi}{6})$ to rectangular co-ordinates
[b] the point with rectangular co-ordinates $(-6, -2)$ to polar co-ordinates
[c] the rectangular equation $x^2 - y^2 - 2x = 0$ to polar
[d] the polar equation $r = \frac{7}{4 - 2\cos\theta}$ to rectangular
[e] the rectangular equation $3x - 2y + 6 = 0$ to polar
[f] the polar equation $r = \cos 2\theta$ to rectangular
[g] the rectangular equation $x^2 + 6y - 9 = 0$ to polar
[h] the polar equation $\theta = \frac{5\pi}{6}$ to rectangular
- [3] Run the standard tests for symmetry for the polar equation $r^3 = 1 - \sin 2\theta$, and state the conclusions.
What is the minimum interval of θ – values that must be plotted before using symmetry to complete the graph ?
- [4] Find the values of $\theta \in [0, 2\pi)$ at which the graph of the polar equation $r = 2\cos 2\theta + 1$ passes through the pole.
- [5] Name the shape of the graphs of the following polar equations.
If the graph is a rose curve, state the number of petals.
- [a] $r = 5 - 5\sin\theta$ [b] $r = 7\sin 6\theta$ [c] $r = 5$ [d] $\theta = 7$
[e] $r = 4\sin 9\theta$ [f] $r = 2 + 3\cos\theta$ [g] $r = 6 - 4\cos\theta$ [h] $r = 3\sin\theta$
[i] $r = 6 + 2\sin\theta$
- [6] Sketch the graphs of the polar equations in [5][a], [f], [g] and [i] using the shortcut process shown in lecture.
Find all x – and y – intercepts.
- [7] Determine if each polar equation corresponds to a circle, a parabola, an ellipse or a hyperbola.
If the equation corresponds to a circle, find its center & radius.
If the equation corresponds to a parabola, find its eccentricity, focus, directrix & vertex.
If the equation corresponds to an ellipse, find its eccentricity, foci, directrix, center & the endpoints of the major axes and latera recta.
If the equation corresponds to a hyperbola, find its eccentricity, foci, directrix, center, vertices & the endpoints of the latera recta.
Do not convert the equations to rectangular co-ordinates.
Final answers must be in rectangular co-ordinates.
- [a] $r = \frac{10}{3 - 3\sin\theta}$ [b] $r = \frac{10}{3 - 2\cos\theta}$ [c] $r = \frac{10}{2 + 3\sin\theta}$ [d] $r = 10$

[2] [a] $(-4\sqrt{3}, 4)$ [b] $(2\sqrt{10}, 3.46)$
 [c] $r = \frac{2\cos\theta}{\cos 2\theta} = 2\cos\theta \sec 2\theta$ [d] $12x^2 + 16y^2 - 28x - 49 = 0$
 [e] $r = \frac{6}{2\sin\theta - 3\cos\theta}$ [f] $(x^2 + y^2)^3 = (x^2 - y^2)^2$
 [g] $r = \frac{3}{1 + \sin\theta}$ or $\frac{-3}{1 - \sin\theta}$ [h] $y = -\frac{\sqrt{3}}{3}x$

[3] Symmetry over polar axis: substituting $(r, -\theta)$ gives $r^3 = 1 + \sin 2\theta$ no conclusion
 substituting $(-r, \pi - \theta)$ gives $r^3 = -1 - \sin 2\theta$ no conclusion
 Symmetry over pole: substituting $(-r, \theta)$ gives $r^3 = -1 + \sin 2\theta$ no conclusion
 substituting $(r, \pi + \theta)$ gives $r^3 = 1 - \sin 2\theta$ symmetric over pole
 Symmetry over $\theta = \frac{\pi}{2}$: substituting $(-r, -\theta)$ gives $r^3 = -1 - \sin 2\theta$ no conclusion
 substituting $(r, \pi - \theta)$ gives $r^3 = 1 + \sin 2\theta$ no conclusion
 Minimum interval $\theta \in [0, \pi]$ or $\theta \in [-\frac{\pi}{2}, \frac{\pi}{2}]$

[4] $0 = 2\cos 2\theta + 1$ for $0 \leq \theta < 2\pi$
 $\cos 2\theta = -\frac{1}{2}$
 $2\theta = \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{8\pi}{3}, \frac{10\pi}{3}$ since $0 \leq 2\theta < 4\pi$
 $\theta = \frac{\pi}{3}, \frac{2\pi}{3}, \frac{4\pi}{3}, \frac{5\pi}{3}$

[5] [a] cardioid [b] rose curve with 12 petals [c] circle [d] line
 [e] rose curve with 9 petals [f] limaçon with inner loop [g] limaçon with dimple [h] circle
 [i] convex limaçon



[7] [a] PARABOLA
 Eccentricity: 1
 Focus: $(0, 0)$
 Directrix: $y = -\frac{10}{3}$
 Vertex: $(0, -\frac{5}{3})$

[b] ELLIPSE
 Eccentricity: $\frac{2}{3}$
 Foci: $(0, 0)$ and $(8, 0)$
 Directrix: $x = -5$
 Center: $(4, 0)$
 Endpoints of major axis: $(-2, 0)$ and $(10, 0)$
 Endpoints of latera recta: $(0, \pm \frac{10}{3})$ and $(8, \pm \frac{10}{3})$

[c] HYPERBOLA

Eccentricity: $\frac{3}{2}$
 Foci: $(0, 0)$ and $(0, 12)$
 Directrix: $y = \frac{10}{3}$
 Center: $(0, 6)$
 Vertices: $(0, 2)$ and $(0, 10)$
 Endpoints of latera recta: $(\pm 5, 0)$ and $(\pm 5, 12)$

[d] Center: $(0, 0)$
 Radius: 10

[8] [a] $r = \frac{7}{1 + \cos \theta}$ [b] $r = \frac{14}{1 - \sin \theta}$ [c] $r = \frac{15}{4 + 3 \sin \theta}$
 [d] $r = \frac{8}{3 - \cos \theta}$ [e] $r = \frac{15}{2 - 5 \cos \theta}$ [f] $r = \frac{15}{2 - 3 \sin \theta}$

[9] [a] ellipse [b] (part of) hyperbola [c] parabola

[10] [a] $y = \frac{-1}{2x + 1}$ [b] $\frac{(y-4)^2}{4} - \frac{(x-3)^2}{25} = 1$ [c] $\frac{(x-8)^2}{36} + (y-7)^2 = 1$

[d] $y = \frac{1}{32} e^{\frac{3x}{5}}$ [e] $y = x^{-\frac{1}{3}}$ [f] $x = \frac{y^2}{2} - 1$

[11] [a] $x = 15t$
 $y = 6 + 15\sqrt{3}t - 16t^2$ [b] over BJ's head

[12] [a] $x = -3 + 10t$
 $y = -6 + 4t$ [b] $x = 2 + \sqrt{29} \cos t$
 $y = -4 + \sqrt{29} \sin t$ [c] $x = t$
 $y = 2t^4 - 3t^3 + 1$
 $t \in [-1, 2]$

[13] All the parametric equations correspond to the line $x = 1 - y$ or $y = 1 - x$

Parametric equations 1:

As t goes from $-\infty$ to ∞ , $y = t^4$ goes from ∞ to 0 to ∞ .

The parametric curve starts in the upper left side of quadrant 2, goes to the x -axis, then goes back to the upper left side of quadrant 2.

Parametric equations 2:

As t goes from $-\infty$ to ∞ , $y = e^t$ goes from 0 to ∞ .

The parametric curve starts near the x -axis, then goes to the upper left side of quadrant 2.

Parametric equations 3:

As t goes from 0 to ∞ , $y = \ln t$ goes from $-\infty$ to ∞ .

The parametric curve starts in the lower right side of quadrant 4, then goes through quadrant 1 to the upper left side of quadrant 2.

Parametric equations 4:

As t goes from $-\infty$ to ∞ , $y = \sin t$ goes back and forth between -1 and 1 .

The parametric curve goes back and forth between the points $(2, -1)$ and $(0, 1)$.

