

## Row Echelon Form (REF)

A matrix is in row echelon form if and only if

the first (leftmost) non-zero entry in each row is 1 (called the leading 1),

the leading 1 in each row (except row 1) is to the right of the leading 1 in the row above it,

and all rows which contain only 0 are below all rows which contain any non-zero entry.

**A matrix in REF corresponds to a system of equations that needs only back-substitution to solve.**

**Are these matrices in REF ? If not, why not ?**

$$\begin{bmatrix} 1 & 3 & 0 & -2 & 4 \\ 0 & 1 & 7 & 4 & 0 \\ 0 & 0 & -1 & 5 & 6 \\ 0 & 0 & 0 & 1 & 3 \end{bmatrix} \quad \begin{bmatrix} 1 & 3 & 0 & -2 & 4 \\ 0 & 1 & 7 & 4 & 0 \\ 0 & 1 & 4 & -3 & -2 \\ 0 & 0 & 1 & 1 & 3 \end{bmatrix} \quad \begin{bmatrix} 1 & 3 & 0 & -2 & 4 \\ 0 & 1 & 7 & 0 & 0 \\ 0 & 0 & 1 & 0 & -2 \\ 0 & 0 & 0 & 1 & 3 \end{bmatrix} \quad \begin{bmatrix} 1 & 3 & 0 & -2 & 4 \\ 0 & 1 & 7 & 4 & 0 \\ 0 & 0 & 0 & 1 & -2 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

## Reduced Row Echelon Form (RREF)

A matrix is in reduced row echelon form if and only if

it is in row echelon form,

and all columns which contain a leading 1 contain only 0 in all other entries.

**A matrix in RREF corresponds to a system of equations that needs the least amount of algebra to solve.**

**Are these matrices in RREF ? If not, why not ?**

$$\begin{bmatrix} 1 & 0 & -1 & -2 & 4 \\ 0 & 1 & 0 & 4 & 0 \\ 0 & 0 & 1 & 5 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 & 0 & 0 & 4 \\ 0 & 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 & -2 \\ 0 & 0 & 0 & 1 & 3 \end{bmatrix} \quad \begin{bmatrix} 1 & 0 & -3 & 0 & 4 \\ 0 & 1 & 8 & 0 & 0 \\ 0 & 0 & 0 & 1 & 6 \\ 0 & 0 & 0 & 0 & 0 \end{bmatrix}$$

---

## Gaussian Elimination Pivot Method

Step 1: Find the first (leftmost) column which contains a non-zero entry

Step 2: Choose a pivot in that column (to be used to replace all lower entries in that column with 0)

Step 3: SWAP to move the pivot's row to the top

Step 4: SCALE to turn the pivot into 1

Step 5: REPLACE each row below the pivot's row

by adding the multiple of the pivot's row which gives a 0 under the pivot

Step 6: Cover up the pivot's row & repeat the entire process (stop when matrix is in row echelon form)

## Gauss-Jordan Elimination (after matrix is in row echelon form)

Step 7: Find the last (rightmost) column which contains a pivot (leading 1)

Step 8: REPLACE each row above the pivot's row

by adding the multiple of the pivot's row which gives a 0 above the pivot

Step 9: Cover up the pivot's row & repeat the entire process (stop when matrix is in reduced row echelon form)

**The following examples should not require fractions if solved using the processes above.**

Example 1:

$$\begin{aligned} 3x + 2y - z &= -1 \\ 5x + y - 3z &= -2 \\ 2x + 4y + 2z &= 2 \end{aligned}$$

Example 2:

$$\begin{aligned} 4x + 6y - 3z &= -15 \\ 3x + 4y + z &= 11 \\ -x - 2y + z &= 1 \end{aligned}$$

Example 3:

$$\begin{aligned} 3x + 4y - 11z &= -17 \\ 2x + y - 4z &= 5 \\ -x - 2y + 5z &= -9 \end{aligned}$$

Example 4:

$$\begin{aligned} 3x + 5y - 9z &= 14 \\ 2x - 3y + 13z &= 3 \\ -x + 2y - 8z &= -1 \end{aligned}$$

Example 5:

$$\begin{aligned} 2x + 4y + 11z &= 10 \\ x + 2y + 7z &= 5 \\ 3x + 4y + 9z &= 13 \end{aligned}$$

Example 1:

CHOOSE 2 in column 1, row 3 as pivot  
(to avoid fractions after scaling)

$$\left[ \begin{array}{ccc|c} 3 & 2 & -1 & -1 \\ 5 & 1 & -3 & -2 \\ \textcircled{2} & 4 & 2 & 2 \end{array} \right] R_1 \leftrightarrow R_3 \Rightarrow \left[ \begin{array}{ccc|c} \textcircled{2} & 4 & 2 & 2 \\ 5 & 1 & -3 & -2 \\ 3 & 2 & -1 & -1 \end{array} \right] R_1 \times \frac{1}{2} \Rightarrow \left[ \begin{array}{ccc|c} \textcircled{1} & 2 & 1 & 1 \\ 5 & 1 & -3 & -2 \\ 3 & 2 & -1 & -1 \end{array} \right] R_2 + (-5)R_1 \Rightarrow R_3 + (-3)R_1$$

SWAP to move pivot to top row      SCALE to turn pivot into 1      REPLACE to eliminate all entries below pivot

COVER row 1 until matrix in REF

CHOOSE -4 in column 2, row 3 as pivot  
(to avoid fractions after scaling)

$$\left[ \begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & -9 & -8 & -7 \\ 0 & \textcircled{-4} & -4 & -4 \end{array} \right] R_2 \leftrightarrow R_3 \Rightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & \textcircled{-4} & -4 & -4 \\ 0 & -9 & -8 & -7 \end{array} \right] R_2 \times \frac{1}{-4} \Rightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & \textcircled{1} & 1 & 1 \\ 0 & -9 & -8 & -7 \end{array} \right] R_3 + (9)R_2$$

SWAP to move pivot to top row      SCALE to turn pivot into 1      REPLACE to eliminate all entries below pivot

COVER row 2 until matrix in REF

CHOOSE 1 in column 3, row 3 as pivot

$$\left[ \begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & \textcircled{1} & 2 \end{array} \right] \Rightarrow \left[ \begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & \textcircled{1} & 2 \end{array} \right] \text{REF}$$

SWAP to move pivot to top row      SCALE to turn pivot into 1

**UNNECESSARY**      **UNNECESSARY**

Rightmost leading 1 in column 3 is pivot

$$\left[ \begin{array}{ccc|c} 1 & 2 & 1 & 1 \\ 0 & 1 & 1 & 1 \\ 0 & 0 & \textcircled{1} & 2 \end{array} \right] R_1 + (-1)R_3 \Rightarrow R_2 + (-1)R_3$$

REPLACE to eliminate all entries above pivot

COVER row 3 until matrix in RREF

Rightmost leading 1 in column 2 is pivot

$$\left[ \begin{array}{ccc|c} 1 & 2 & 0 & -1 \\ 0 & \textcircled{1} & 0 & -1 \\ 0 & 0 & 1 & 2 \end{array} \right] R_1 + (-2)R_2$$

REPLACE to eliminate all entries above pivot

COVER row 2 until matrix in RREF

$$\left[ \begin{array}{ccc|c} 1 & 0 & 0 & 1 \\ 0 & 1 & 0 & -1 \\ 0 & 0 & 1 & 2 \end{array} \right] \text{RREF} \Rightarrow \begin{array}{l} x = 1 \\ y = -1 \\ z = 2 \end{array}$$

CHECK:

$$\begin{array}{l} 3(1) + 2(-1) - (2) = 3 - 2 - 2 = -1 \\ 5(1) + (-1) - 3(2) = 5 - 1 - 6 = -2 \\ 2(1) + 4(-1) + 2(2) = 2 - 4 + 4 = 2 \end{array}$$