



Thermodynamics

Heat Capacity

Phase Changes

Lana Sheridan

De Anza College

April 24, 2018

Last time

- finish applying the ideal gas equation
- thermal energy
- introduced heat capacity

Overview

- heat capacity
- phase changes
- latent heat

Heat Capacity

Before we look more closely at the first law, let's look at the effect of adding heat to a substance that is not near a phase change.

It requires less energy to raise the temperature of some objects compared to others.

Heat Capacity

Before we look more closely at the first law, let's look at the effect of adding heat to a substance that is not near a phase change.

It requires less energy to raise the temperature of some objects compared to others.

Obviously, a small amount of water requires less heat to raise its temperature by 1 degree than a large amount of water.

But even two objects of the same mass may require different amounts of heat to change their temperature by 1 degree if they are made of different materials.

Different materials have different **heat capacities**.

Heat Capacity

Heat Capacity, C

of a sample of substance is the quantity of heat required to change the temperature **of that sample** by 1 degree C (or K).

$$Q = C \Delta T$$

where ΔT is the change in temperature and Q is the heat.

Specific Heat Capacity, c

of a substance is the quantity of heat required to change the temperature of a unit mass of that substance by 1 degree C (or K).

$$Q = cm \Delta T$$

m is the mass of the object.

Specific Heat Capacity

Different materials have different heat capacities.

- for Lead, $c = 129 \text{ J kg}^{-1} \text{ K}^{-1}$
- for Hydrogen, $c = 14300 \text{ J kg}^{-1} \text{ K}^{-1}$

Hydrogen gas's heat capacity is phenomenally high. (Its molar mass is small.)

Most substances are in the range $500 - 2000 \text{ J kg}^{-1} \text{ K}^{-1}$.

This means that water also has quite a high heat capacity ($4186 \text{ J kg}^{-1} \text{ K}^{-1}$). This has an effect on Earth's weather and climate, since oceans make most of Earth's surface.

Specific Heat Capacity Question

Quick Quiz 20.1¹ Imagine you have 1 kg each of iron, glass, and water, and all three samples are at 10°C.

(a) Rank the samples from highest to lowest temperature after 100 J of energy is added to each sample.

Heat capacities: glass – 837 J kg⁻¹ K⁻¹

iron – 448 J kg⁻¹ K⁻¹

- (A) iron, glass, water
- (B) water, iron, glass
- (C) water, glass, iron
- (D) glass, iron, water

¹Serway & Jewett, pg 579.

Specific Heat Capacity Question

Quick Quiz 20.1¹ Imagine you have 1 kg each of iron, glass, and water, and all three samples are at 10°C .

(b) Rank the samples from greatest to least amount of energy transferred by heat if each sample increases in temperature by 20°C .

- (A) iron, glass, water
- (B) water, iron, glass
- (C) water, glass, iron
- (D) glass, iron, water

¹Serway & Jewett, pg 579.

Heat and Temperature Change

Energy that causes a change in temperature does not have to enter our system as heat.

It can be a different form of energy transfer.

Examples:

- in a microwave, energy T_{ER} enters the food as **electromagnetic waves**
- **work** can cause a temperature change in two surfaces rubbed together, or as a bicycle pump pressurizes air in the bike tires, the air's temperature rises

These energy transfers to our system will increase the internal energy of the system, E_{int} .

Calorimetry


Calorimetry

a technique for determining the specific heat capacity of a sample by heating it to a known temperature, then transferring it to a known quantity of water and observing the temperature change in the water.

Steps:

- 1 sample of known mass m_x is heated to temperature T_x
- 2 sample is moved to an isolated container of water, containing mass m_w of water at temperature $T_w < T_x$
- 3 the sample and the water are allowed to reach thermal equilibrium
- 4 the final temperature of the water, T_f , is measured

Calorimetry



Since the heat transferred to the cold water is equal to the heat transferred from the hot sample:

$$Q_c = -Q_h$$
$$m_w c_w (T_f - T_w) = -m_x c_x (T_f - T_x)$$
$$c_x = \frac{m_w c_w (T_f - T_w)}{m_x (T_x - T_f)}$$

¹Figure from Serway & Jewett, page 595.

Phase Changes

The processes by which matter changes from one state to another.


The different states of matter: solid, liquid, gas, plasma, are also called *phases* of matter.

Phase Changes

Phase changes tend to be dramatic.

If sudden, obvious changes in the properties and behaviors of a substance did not occur as we vary the temperature, we would probably have no need to refer to different states of matter!

Phase Changes




Notice the discontinuity!

Phase Changes

We know that as we heat a solid it will eventually melt to form a liquid and if we keep heating the liquid will boil off as a gas.

But how does the temperature change during these processes?

Phase Changes



During a phase change, temperature doesn't change, even when heat is added!

Phase Changes

Why does this happen?

Where is the energy going?

It isn't increasing the translational speed of the atoms, that would relate to an increase in temperature.

Latent Heat

latent heat of fusion, L_f

The amount of energy (heat) per unit mass required to change a solid to a liquid.

$$Q = mL_f$$

where m is the mass of solid that is transformed into a liquid.

latent heat of vaporization, L_v

The amount of energy (heat) per unit mass required to change a liquid to a gas.

$$Q = mL_v$$

where m is the mass of liquid that is transformed into a gas.

¹“Latent” from *latere*, “to lie hidden”.

Latent Heat

Substance	Melting Point (°C)	Latent Heat of Fusion (J/kg)	Boiling Point (°C)	Latent Heat of Vaporization (J/kg)
Helium ^a	-272.2	5.23×10^3	-268.93	2.09×10^4
Oxygen	-218.79	1.38×10^4	-182.97	2.13×10^5
Nitrogen	-209.97	2.55×10^4	-195.81	2.01×10^5
Ethyl alcohol	-114	1.04×10^5	78	8.54×10^5
Water	0.00	3.33×10^5	100.00	2.26×10^6
Sulfur	119	3.81×10^4	444.60	3.26×10^5
Lead	327.3	2.45×10^4	1 750	8.70×10^5
Aluminum	660	3.97×10^5	2 450	1.14×10^7
Silver	960.80	8.82×10^4	2 193	2.33×10^6
Gold	1 063.00	6.44×10^4	2 660	1.58×10^6
Copper	1 083	1.34×10^5	1 187	5.06×10^6

^aHelium does not solidify at atmospheric pressure. The melting point given here corresponds to a pressure of 2.5 MPa.

Latent Heat

Latent heat is the energy required for the to change to the higher energy phase per unit mass of the substance.

This includes two components:

- 1 the energy required to overcome intermolecular forces / break the bonds
- 2 the work required to push aside gas at ambient pressure to allow for any increased volume of the new phase

¹This is relevant in Ch 20, problem 77.

Latent Heat

Latent heat is the energy required for the to change to the higher energy phase per unit mass of the substance.

This includes two components:

- 1 the energy required to overcome intermolecular forces / break the bonds
- 2 the work required to push aside gas at ambient pressure to allow for any increased volume of the new phase

The latent heat depends on the temperature and pressure of the phase change.

¹This is relevant in Ch 20, problem 77.

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Supposing that it remains at that value all the way to absolute zero, calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water. How does this number of calories required to change the same gram of 100°C boiling water to 100°C steam?

Reminder: 1 cal is the heat required to raise the temperature of 1 g of water by 1°C .

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water.

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water.

warming ice:

$$Q_1 = m_{\text{ice}} \Delta T = (1 \text{ g})(0.5 \text{ cal/g}^\circ\text{C})(273^\circ\text{C}) = 136.5 \text{ cal}$$

melting:

$$Q_2 = mL_f = (1 \text{ g}) \left(\frac{3.33 \times 10^5 \text{ J/kg}}{4.186 \text{ J/cal}} \right) \left(\frac{1 \text{ kg}}{1000 \text{ g}} \right) = 79.55 \text{ cal}$$

warming water:

$$Q_3 = m_{\text{water}} \Delta T = (1 \text{ g})(1.0 \text{ cal/g}^\circ\text{C})(100^\circ\text{C}) = 100 \text{ cal}$$

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water.

warming ice:

$$Q_1 = m_{\text{ice}} \Delta T = (1 \text{ g})(0.5 \text{ cal/g}^\circ\text{C})(273^\circ\text{C}) = 136.5 \text{ cal}$$

melting:

$$Q_2 = mL_f = (1 \text{ g}) \left(\frac{3.33 \times 10^5 \text{ J/kg}}{4.186 \text{ J/cal}} \right) \left(\frac{1 \text{ kg}}{1000 \text{ g}} \right) = 79.55 \text{ cal}$$

warming water:

$$Q_3 = m_{\text{water}} \Delta T = (1 \text{ g})(1.0 \text{ cal/g}^\circ\text{C})(100^\circ\text{C}) = 100 \text{ cal}$$

Total $Q_1 + Q_2 + Q_3 = 320 \text{ cal}$.

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Supposing that it remains at that value all the way to absolute zero, calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water. How does this number of calories required to change the same gram of 100°C boiling water to 100°C steam?

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Supposing that it remains at that value all the way to absolute zero, calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water. How does this number of calories required to change the same gram of 100°C boiling water to 100°C steam?

boiling:

$$Q_4 = mL_v = (1 \text{ g}) \left(\frac{2.26 \times 10^6 \text{ J/kg}}{4.186 \text{ J/cal}} \right) \left(\frac{1 \text{ kg}}{1000 \text{ g}} \right) = 540 \text{ cal}$$

Practice

The specific heat capacity of ice is about $0.5 \text{ cal/g}^\circ\text{C}$. Supposing that it remains at that value all the way to absolute zero, calculate the number of calories it would take to change a 1 g ice cube at absolute zero (-273°C) to 1 g of boiling water. How does this number of calories required to change the same gram of 100°C boiling water to 100°C steam?

boiling:

$$Q_4 = mL_v = (1 \text{ g}) \left(\frac{2.26 \times 10^6 \text{ J/kg}}{4.186 \text{ J/cal}} \right) \left(\frac{1 \text{ kg}}{1000 \text{ g}} \right) = 540 \text{ cal}$$

The energy required to transform the water to steam is much bigger than the energy required to heat the ice, convert it to water, and continue heating up to 100°C .

Question

Suppose the same process of adding energy to the ice cube is performed as discussed in the last question, but instead we graph the internal energy of the system as a function of energy input. What would this graph look like?

¹Based on Quick Quiz 20.2, Serway & Jewett, page 600.

Practice

The heat of vaporizations of ethyl alcohol is about 200 cal/g. If 2 kg of this fluid were allowed to vaporize in a refrigerator, show that 5 kg of ice (at 0°C) would be formed from 0°C water.

Practice

The heat of vaporizations of ethyl alcohol is about 200 cal/g. If 2 kg of this fluid were allowed to vaporize in a refrigerator, show that 5 kg of ice (at 0°C) would be formed from 0°C water.

Hint: in the last problem we melted 1 g of ice and found it required 80 cal.

Practice

The heat of vaporizations of ethyl alcohol is about 200 cal/g. If 2 kg of this fluid were allowed to vaporize in a refrigerator, show that 5 kg of ice (at 0°C) would be formed from 0°C water.

Hint: in the last problem we melted 1 g of ice and found it required 80 cal.


energy needed for vaporization:

$$Q = mL_{v,ea} = (2 \text{ kg})(200 \text{ cal/g}) = 4.0 \times 10^5 \text{ cal}$$

assuming this same amount of energy was taken from the water:

$$m = \frac{Q}{L_f} = \frac{4.0 \times 10^5 \text{ cal}}{80 \text{ cal/g}} = 5000 \text{ g} = 5 \text{ kg} \quad \checkmark$$

Phase Change paths



Evaporation

evaporation

the process by which a liquid changes to a gas at the liquid surface

Since changing from a liquid to a gas requires heat, when a liquid evaporates it takes heat from its surroundings.

This is why humans sweat in hot weather, pigs wallow puddles, and dogs pant. All are trying to use evaporation of water to reduce body temperature.

Evaporation

Ben Franklin noticed that a wet shirt kept him feeling cool on a hot day.


He decided to experiment to see if the temperature of objects could be lowered by this process.

In 1758 he and John Hadley took a mercury thermometer and repeatedly wet the bulb with ether while using bellows to keep air moving over it.

Despite it being a warm day, they recorded temperatures as low as 7°F (-14°C) at the bulb of the thermometer.

This is the basic idea behind refrigeration!

Phase Diagrams



¹A typical phase diagram. The dashed green line shows the unusual behavior of water. Diagram by Matthieumarechal, Wikipedia.

Summary

- heat capacity
- phase changes
- latent heat

Homework Serway & Jewett:

- Look at examples 20.1–4.
- prev: Ch 20, onward from page 615. OQs: 3, 5, 7; CQs: 3, 11; Probs: 1, 3, 9, 13, 63, 69
- new: Ch 20. Probs: 19, 23, 71, 77