

1. State the 3 important parts of the Central Limit Theorem for the sample mean  $\bar{X}$

**Mean stays the same**

**Standard deviation gets smaller** in proportion to the square root of the sample size

When  $n$  is sufficiently large, sample mean has approximately a **Normal Distribution**

2. The completion time (in minutes) for a student to complete a short quiz follows the continuous probability density function shown here, with some areas calculated. It is known that  $\mu=5.3$  minutes and  $\sigma = 2.4$  minutes. 40 students take the quiz.

- a. Find the probability that the **mean** completion time for the students is under 5 minutes.

$$P(X < 5) = P\left(Z < \frac{5-5.3}{2.4/\sqrt{40}}\right) = P(Z < -0.79) = 0.2148$$

- b. Find the probability that the **mean** time for the class to finish the quiz is between 5 and 6 minutes.

$$P(6 \leq X \leq 8) = P\left(\frac{5-5.3}{2.4/\sqrt{40}} \leq Z \leq \frac{6-5.3}{2.4/\sqrt{40}}\right) = P(-0.79 \leq Z \leq 1.84) = 0.9671 - 0.2148 = 0.7523$$

- c. The **mean** completion time for the class was 7.1 minutes. Is this result unusual? Explain.

$$Z = \frac{7.1-5.3}{2.4/\sqrt{40}} = 4.74 \text{ Yes, this is a very unusual result}$$

3. For women aged 18-24, systolic blood pressures (in mmHg) are normally distributed with  $\mu=114.8$  and  $\sigma=13.1$ .

- a. Find the probability a woman aged 18-24 has systolic blood pressure exceeding 120.

$$P(X > 120) = P\left(Z > \frac{120-114.8}{13.1}\right) = P(Z > 0.40) = 1 - 0.6554 = 0.3446$$

- b. If 40 women are randomly selected, find the probability that their mean blood pressure exceeds 120.

$$P(\bar{X} > 120) = P\left(Z > \frac{120-114.8}{13.1/\sqrt{40}}\right) = P(Z > 2.51) = 1 - 0.9940 = 0.0060$$

- c. If the pdf for systolic blood pressure did NOT follow a normal distribution, would your answer to part b change?

**Answer would stay the same. When  $n$  is sufficiently large, Sample mean has approximately a Normal Distribution regardless of the shape of the pdf of  $X$ .**

4. It has been reported that 28% of all Californians have visited Yosemite National Park. A pollster samples 1000 Californians and determines that 240 of them have visited Yosemite National Park.
- a. Determine the value of the sample proportion,  $\hat{p}$ , of Californians who have visited Yosemite. Is the higher or lower than the reported population value?

$$\hat{p} = 240 / 1000 = 0.24 \text{ This is lower than the reported population value}$$

- b. Determine the expected value of the sample proportion.

$$\mu_{\hat{p}} = p = 0.28$$

- c. Determine the standard deviation of the sample proportion.

$$\sigma_{\hat{p}} = \sqrt{\frac{0.28(1-0.28)}{1000}} = 0.0142$$

- d. Determine that the condition for normality is satisfied.

$$1000(.28)=280 \quad 1000(1-.28) = 720. \text{ Both values are greater than 10, so conditions for normality are met.}$$

- e. Determine the probability the sample proportion exceeds 0.24.

$$P(\hat{p} > 0.24) = P\left(Z > \frac{0.24 - 0.28}{0.0142}\right) = P(Z > -2.82) = 1 - 0.0024 = 0.9976$$

- f. Determine the probability the sample proportion is between 0.2 and 0.3.

$$P(0.2 \leq \hat{p} \leq 0.3) = P\left(\frac{0.2 - 0.28}{0.0142} \leq Z \leq \frac{0.3 - 0.28}{0.0142}\right) = P(-5.63 \leq Z \leq 1.41) = 0.9207 - 0 = 0.9207$$